

---

# TRINITY FLUIDS, LLC

Designed for Quality, Value and Professionals

---

**SaniSol 7** is formulated to protect the employee from many unseen germs, bacteria and viruses found in the morgues, prep rooms, labs and restrooms.

## Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SaniSol7 is not for use on critical or semi-critical medical device surfaces.

Fill or refill the SaniSol 7 spray bottle with SaniSol 7 refill. Pour product into spray bottle over sink or tub basin.

**DISINFECTION, DEODORIZING AND CLEANING** – Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for **3 minutes** (for Blood borne Pathogens: allow surface to remain wet for HIV-1 **1 minute** and for HBV and HCV **5 minutes**). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. SaniSol7 will not leave grit or soap scum.

Using approved AOAC test methods, in the presence of 5% blood serum and a 3 minute contact time, unless otherwise noted, SaniSol7 kills the following organism on hard non-porous inanimate surfaces:

### Bactericidal Activity (Bacteria):

Pseudomonas aeruginosa, (ATCC 15442)	Listeria monocytogenes, (ATCC 35152)
Staphylococcus aureus, (ATCC 6538)	Yersinia enterocolitica, (ATCC 23715)
Salmonella (choleraesuis) enteric, (ATCC 10708)	Enterococcus faecium (ATCC 6569)
Escherichia coli, (ATCC 11229)	Corynebacterium ammoniagenes, (ATCC 6871)
Escherichia coli O157:H7, (ATCC 43895)	Salmonella (typhi) enteric, (ATCC 6539)
Streptococcus pyogenes (Necrotizing Fasciitis-Group A) (V.A. Medical Center Isolate 04001)	

### Antibiotic-Resistant (Strains of)(Antibiotic-Resistant Bactericidal Activity):

Methicillin resistant Staphylococcus aureus (MRSA), (ATCC 33593)  
Vancomycin resistant Enterococcus faecalis (VRE), (ATCC 51575)  
Vancomycin intermediate resistant Staphylococcus aureus (VISA), CDC Isolate 99287)  
Methicillin resistant Staphylococcus epidermidis (MRSE), (ATCC 51625)  
Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA), (NRS 123) Genotype USA400  
Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA), (NRS 384) Genotype USA300

**Tuberculocidal Activity** – SaniSol 7 exhibits disinfectant efficacy against Mycobacterium tuberculosis BCG (Mycobacterium Bovis) at 20 degrees Centigrade when the treated surface is allowed to remain wet for **5 minutes**.

**\*Virucidal Activity** – SaniSol 7 kills on hard, non-porous inanimate surfaces when allowed to remain wet for a **10 minute** contact time against:

Canine Parvovirus (ATCC VR-2017)  
Hepatitis A Virus (HAV) (University of Ottawa)  
Poliovirus Type 1(ATCC VR-1000)

SaniSol7, when used on environmental, inanimate hard surfaces exhibits Virucidal activity against: **Hepatitis B Virus (HBV)**(Hepadnavirus Testing, Inc.), **Hepatitis C Virus (HCV)** and **Bovine Viral Diarrhea Virus (BVDV)** (NVSL) when the treated surface is allowed to remain wet for **5 minutes**; **Human Coronavirus** (ATCC VR-740), **SARS associated Coronavirus** (ZeptoMetrix), **Avian Influenza A strain H3N2** (ATCC VR-2072) and **Avian Influenza A strain H9N2** (Charles River Laboratories) (Avian Influenza strains H3N2 and H9N2) for **2 minutes**; **Paramyxovirus (Mumps)** (ATCC

VR-1438), **Rhinovirus type 39** (ATCC VR-340) and **Rotavirus** (University of Ottawa) for **3 minutes**; **HIV-1 (associated with AIDS)** (Advanced Biotechnologies, Inc) for **1 minute (60 seconds)**; and **Norovirus (Norwalk virus)**, **Feline Calicivirus** (University of Ottawa) and **Rabies Virus** (CDC) for **30 seconds**.

**KILLS HIV-1, HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

**PERSONAL PROTECTION:** When handling items soiled with blood or body fluids use the disposable latex gloves, gowns masks or eye coverings

**CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

**CONTACT TIME:** Allow surface to remain wet for HBV and HCV 5 minutes, for HIV-1 1 minute. These contact times will not control other common types of viruses and bacteria.

**DISPOSAL OF INFECTIOUS MATERIAL:** Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

### **Precautionary Statements**

#### **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

##### **CAUTION**

Keep out of reach of Children. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

##### **First Aid**

Have product container or label with you when calling a poison central or doctor or going for treatment.

**If in eyes:** Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance then give CPR, preferably by mouth to mouth, if possible. Call a poison control center or doctor for further treatment advice.

##### **Note to Physician**

Probable mucosal damage may contraindicate the use of gastric lavage.

(Product supplied in containers 5 gallons or greater)

##### **Environmental Hazards**

This pesticide is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying sewage treatment plant authority.

#### **Storage and Disposal**

**DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.**

Store in a dry place no lower in temperature than 50 degree F. or higher than 120 degree F. Do not reuse empty container (Cubetainer).

*Container Disposal for Refill Spray Bottle:* Do not reuse empty spray bottle except with refill Cubetainer, otherwise discard in trash. Do not reuse empty Cubetainer. Breakdown empty Cubetainer and discard in trash or recycle per local and state regulations.